

Supplementary Table 1

| | Age (y) | Gender | Oral ulcer score | Genital ulcer score | Acne-like lesion score | Erythema nodosum-like lesion score | NRS of pain | Total score | Evaluation |
|--------|---------|--------|------------------|---------------------|------------------------|------------------------------------|-------------|-------------|------------|
| Case 1 | 31 | F | 4.5 | 0 | 4 | 8 | 7 | 23.5 | Moderate |
| Case 2 | 59 | F | 9 | 0 | 5 | 0 | 5 | 19 | Moderate |
| Case 3 | 29 | F | 0 | 0 | 2 | 9 | 9 | 20 | Moderate |
| Case 4 | 59 | F | 6.5 | 0 | 0 | 7.75 | 8 | 22.25 | Moderate |
| Case 5 | 63 | F | 5.5 | 0 | 0 | 7 | 6 | 18.5 | Moderate |
| Case 6 | 49 | F | 7 | 0 | 0 | 0 | 8 | 15 | Moderate |
| Case 7 | 36 | F | 2 | 0 | 0 | 0 | 4 | 6 | Mild |
| Case 8 | 24 | F | 0 | 0 | 1.5 | 0 | 0 | 1.5 | Mild |
| Case 9 | 33 | F | 0 | 0 | 1.5 | 0 | 0 | 1.5 | Mild |

Supplementary Figure 1

